## Analysis of Lake Evaporation at Lake Tahoe, California/Nevada, USA



Healey, N.C.<sup>1</sup>, Hook, S.J.<sup>1</sup>, Lenters, J.D.<sup>2,3,4</sup>, Soylu, M.E.<sup>5</sup>, Schladow, S.G.<sup>6</sup>

<sup>1</sup> NASA Jet Propulsion Laboratory, California, Institute of Technology <sup>2</sup> LimnoTech, Inc.

<sup>3</sup> University of Colorado, Boulder

<sup>4</sup> University of Wisconsin, Madison

<sup>5</sup> Meliksah University, Turkey

<sup>6</sup> University of California at Davis













#### **Outline**



### **Study Site Details**

#### **□** Details

Elevation: 1897 m
Max. Depth: 501 m
Avg. Depth: 300 m
Surface Area: 490 km²

• Climate: Dry-summer continental (Koeppen: Dsb)

- Oligomictic (avg. mixing every ~4 years)
- ~69 inlets, only 1 outlet (Truckee River)


#### **□** Issues

- ✓ Water Quality and Clarity
- ✓ Ecosystem Sustainability
- ✓ Tourism
- ✓ Economics
- ✓ Outdoor Recreation



#### □ Facts

- ➤ Largest alpine lake in North America
- Second deepest lake in the United States
- Sixth largest by volume behind the Great Lakes





Recently, reservoirs across the western United States have been experiencing extremely low water levels with water **demands increasing and supplies decreasing**.

(Fulp, 2005; Barnett and Pierce, 2008)






















Recently, reservoirs across the western United States have been experiencing extremely low water levels with water demands increasing and supplies decreasing. (Fulp, 2005; Barnett and Pierce, 2008)

Open water evaporation is one of the most difficult surface energy/water fluxes to quantify, and is rarely directly measured in the natural environment.

Lake Powell, a reservoir on the Colorado River. Photo


University of Colorado Boulder





Duration of time when lake is at maximum capacity (i.e. Legal Storage Limit) has been decreasing since ~1940.





What are the dominant factors controlling water level at Lake Tahoe?

How influential is evaporation in the overall water budget of Lake Tahoe?






## **Methodology: Estimating Evaporation**

- **□** Pan Evaporation
  - Pan coefficient
- ☐ Bowen Ratio Energy Balance (BREB)
  - Net radiation, heat storage, inflow, outflow, humidity & temperature gradients
- ☐ Eddy Covariance
  - 3-D wind and water vapor
- Water Balance
  - inflow, outflow, groundwater in and groundwater out, diversions, and precipitation
- Models
  - Example: Complimentary Relationship Lake Evaporation (CRLE)
- ☐ Mass Transfer/Bulk Aerodynamic
  - Surface temperature, humidity, and wind speed over water

Water vapor flux from an open water body surface by way of turbulent diffusion as a function of **wind speed** and the **vapor pressure gradient** (Dalton, 1802)

Benefits: Low Cost, Minimal Data Requirements









## Methodology: Buoys at Lake Tahoe

- ☐ Deployed in 1999
- ☐ Continuous
  Measurements
  (every 5-min)



**Custom JPL Radiometers** 


- **□** 8-14 μm
- $\Box$  Accuracy:  $\pm 0.08$  °C
- ☐ Calibrated at JPL with NIST-traceable stirred water bath blackbody







### **Evaporation Estimation at Lake Tahoe: "Hybrid Method"**

Bowen Ratio Energy Balance (BREB) and Mass Transfer



16 different time periods (42-391 days) for BREB analysis using 2011-2012 data





#### "Hybrid Method"

BREB estimates of *H* generate estimates of *LE* using the Bowen Ratio.

$$\beta = \gamma \frac{\overline{U(T_s - T_a)}}{\overline{U(e_s - e_a)}}$$

$$\beta = \frac{H}{LE}$$

#### **Mass Transfer**

$$H = C_H \cdot U(T_s - T_a)$$

$$H = C_H \cdot U(T_s - T_a)$$

$$LE = C_E \cdot U(e_s - e_a)$$

γ : psychrometric constant (m s<sup>-1</sup>)

 $\boldsymbol{U}$ : wind speed (m s<sup>-1</sup>)

 $e_s$ : saturation vapor pressure of water (kPa)


 $e_a$ : vapor pressure of air (kPa)

 $T_s$ : surface water temperature (°C)


 $T_a$ : air temperature (°C)

### **Methodology: Water Balance**

$$(PPT + SW_{in} + GW_{in}) - (Evap + SW_{out} + GW_{out}) = 0$$




#### Assume groundwater exchange is negligible



**Precipitation** data is from the National Weather Service COOP station at Tahoe City


**Inflow** and **outflow** data are from all USGS gaging stations around Lake Tahoe.




### **Methodology: Water Balance**

$$(PPT + SW_{in} + GW_{in}) - (Evap + SW_{out} + GW_{out}) = 0$$

$$(PPT + SW_{in}) - (Evap + SW_{out}) = 0$$




Assume groundwater exchange is negligible



**Precipitation** data is from the National Weather Service COOP station at Tahoe City

**Inflow** and **outflow** data are from all USGS gaging stations around Lake Tahoe.



Results and Discussion: Diel and Daily Evaporation


- Computed rates at Lake Tahoe are comparable with other **alpine lakes** around the world.
  - ightharpoonup Range:  $0.9 6.3 \text{ mm day}^{-1}$


(dos Reis and Dias, 1998; Elsawwaf et al., 2010; Ikebuchi et al., 1988; Gibson et al., 1996; Gibson, 2002; Sene at al., 1991; Saxena, 1996; Kittel and Richerson, 1978; Patrick and Kauahikaua, 2015; Roy and Hayashi, 2008)


- Other estimates of Evaporation at Lake Tahoe:
  - ightharpoonup Range: 2.5 3.0 mm day-1


(Dugan and McGauhey, 1974; Myrup et al., 1979; Trask, 2007; Huntington and McEvoy, 2011)

#### Diel Cycle of Evaporation (2002 – 2015) Lake Tahoe, CA/NV












© 2017 California Institute of Technology, Government sponsorship acknowledged

## Results and Discussion: Daily and Monthly Evaporation




Due to the high thermal inertia of water, peak evaporation occurs between August and October; lagging behind peak solar radiation and air temperature.







## **Results and Discussion: Seasonal Evaporation**

# Winter = Onshore Flow, Wet (Increasing Precipitation) = Water Level Rises












## **Results and Discussion: Seasonal Evaporation**


Winter = Onshore Flow, Wet (Increasing Precipitation) = Water Level Rises



Summer = Offshore Flow, Dry (Increasing Evaporation) = Water Level Falls









## Results and Discussion: Water Level, Precipitation, Evaporation


Precipitation patterns at Lake Tahoe are seasonally wet in winter, and dry in summer.


Water levels dropped below the natural rim in 2005, 2010, 2014-2015.




Anomalously high **precipitation** leads to **higher** lake levels


Anomalously high **evaporation** leads to **lower** lake levels.





#### **Results and Discussion: Annual Water Balance**






Note: Water added (P and SWin) are shown as positive values, and water removed from the lake (E and SWout) are shown as negative values.






#### **Results and Discussion: Annual Water Balance**

Calculated water level change based on water balance: Very close agreement with observations, with a few exceptions






#### Potential Reasons for discrepancies in Calculated Water Level Change:

- Inflow measurement error
   ~69 inlets and only 10 USGS gages
- Precipitation measurement error
- Evaporation estimation error


- Diversions for municipalities ex. South Lake Tahoe, Tahoe City, etc.
- Diversions for commercial uses ex. Golf courses

#### **Results and Discussion: Stratification**



Longer stratification seasons = warmer lake water surface temperatures = increased evaporation


## **Results and Discussion: Lake Ecology**




#### What we can expect:


- Changes in fish species composition.
- Less dissolved organic carbon and oxygen – more visible light and more UV radiation penetration.
- More weed growth – blue green algae growth.
- Longer stratification will alter mixing dynamics

### **Recent Drought Relief**









January 2016

**January 9, 2017** 

#### **Conclusions**

In western USA: demand for water is increasing and supply is decreasing


Need for improved water balance estimates

#### What are the dominant factors controlling water level at Lake Tahoe?

- Precipitation and Evaporation

#### How influential is evaporation in the overall water budget of Lake Tahoe?

- Accounts for roughly 80% of water losses.
- Average annual water lost to evaporation:
  - **❖** 535.8 million m<sup>3</sup>
  - **434,000** acre ft.
  - **❖** 141.5 billion gallons = ~800 thousand households





#### References

- Alpine Space Programme, 2014. SILMAS Alpine Lakes Network: Climate Change Impacts on Alpine Lakes. Alpine Space Programme, EU transnational cooperation programme for the Alps Report. 115 pp. Retrieved from the internet 2/8/2017 at http://www.alpine-space.org/2007-2013/uploads/tx\_txrunningprojects/SILMAS\_Climate\_Change\_\_Impacts\_on\_Alpine\_Lakes\_part1.pdf
- Barnett, T.P., and D.W. Pierce, 2008. When will Lake Mead go dry? Water Resources Research, 44, W03201.
- Dalton, J., 1802. Experimental essays on the constitution of mixed gases: on the force of steam or vapour from water or other liquids in different temperatures, both in a Torricelli vacuum and in air; on evaporation; and on expansion of gases by heat. *Manchester Literary and Philosophical Society Member Proceedings*, 5, 536-602.
- Dugan, G.L., and McGauhey, P.H., 1974. Enrichment of Surface Waters. Journal WPCF, 46(10).
- dos Reis, R.J. and Dias, N.L. 1998. Multi-season lake evaporation: energy-budget estimates and CRLE model assessment with limited meteorological observations. *Journal of Hydrology*, 208, 135-147.
- Elsawwaf, M., *et al.*, 2010. Evaporation estimates from Nasser Lake, Egypt, based on three floating station data and Bowen ratio energy budget. *Theoretical and Applied Climatology*, 100(3), 439-465.
- Fulp, T., 2005. Response of the System to Various Hydrological and Operational Assumptions: Reclamation Modeling Results. University of Colorado Natural Resources Law Center, Twenty-sixth Annual Conference.
- Gibson, J.J., et al., 1996. Development and validation of an isotopic method for estimating lake evaporation. Hydrological Processes, 10, 1369-1382.
- Gibson, J.J, 2002. Short-term evaporation and water budget comparisons in shallow Arctic lakes using non-steady isotope mass balance. *Journal of Hydrology*, 264, 242-261.
- Huntington, J.L., and D. McEvoy, 2011. Climatological estimates of open water evaporation from selected Truckee and Carson River basin water bodies, California and Nevada. Division of Hydrologic Sciences, Desert Research Institute, Reno, NV Publication No. 41254, 1-43.
- Ikebuchi, S., et al., 1988. Evaporation from Lake Biwa. Journal of Hydrology, 427-449.
- Kittel, T., and P.J. Richerson, 1978. The heat budget of a large tropical lake, Lake Titicaca (Peru-Bolivia). Verh. Internat. Verein. Limnol., 20, 1203-1209.
- Lenters, J.D., and M.E. Soylu, 2014. Development of real-time evaporation and energy balance algorithms for Lake Tahoe using in-situ buoy data. LimnoTech, Inc. Technical Report.
- Lenters, J.D., *et al.*, 2005. Effects of Climate variability on lake evaporation: Results from a long-term energy budget study of Sparkling Lake, northern Wisconsin (USA). *Journal of Hydrology*, 308(1-4), 168-195.
- Myrup, L.O., et al., 1979. Climatological estimate of the average monthly energy and water budgets of Lake Tahoe, California-Nevada. *Water Resources Research*, 15, 6, 1499–1508.
- Patrick, M.R., and J. Kauahikaua, 2015. USGS Satellite Monitoring of Dramatic Changes at Hawai'i's only Alpine Lake Lake Waiau on Mauna Kea Volcano. USGS Investigations Report 2015-5076.
- Roy, J.W., and M. Hayashi, 2008. Groundwater exchange with two small alpine lakes in the Canadian Rockies. *Hydrological Processes*, 22, 2838-2846.
- Sahoo, G.B., et al., 2016. Climate change impacts on lake thermal dynamics and ecosystem vulnerabilities. Limnology and Oceanography, 61:496–507.
- Saxena, R.K., 1996. Estimation of lake evaporation from a shallow lake in central Sweden by oxygen-18. *Hydrological Processes*, 10, 1273-1281.
- Sene, K.J., et al., 1991. Evaporation from a tropical lake: comparison of theory with direct measurements. Journal of Hydrology, 127, 193-217.
- Tahoe Environmental Research Center (TERC), University of California Davis (2015) Tahoe: State of the Lake Report 2015. Accessed in June 2016: http://terc.ucdavis.edu/stateofthelake/sotl-reports/2015/complete-2015-sotl\_low.pdf
- Trask, J.C., 2007. Resolving Hydrologic Water Balances through Novel Error Analysis with Focus on Inter-annual and long-term Variability in the Tahoe Basin. University of California, Davis, Ph.D. Dissertation. 378 p.





Contact me at:
<a href="mailto:Nathan.C.Healey@jpl.nasa.gov">Nathan.C.Healey@jpl.nasa.gov</a>

4800 Oak Grove Dr. Pasadena, CA 91109 818.393.5358

